

# ORIENTING GRAPHS TO OPTIMIZE REACHABILITY\*

S. Louis Hakimi,<sup>†</sup> Edward F. Schmeichel,<sup>‡</sup> Neal E. Young<sup>§</sup>

## Abstract

It is well known that every 2-edge-connected graph can be oriented so that the resulting digraph is strongly connected. Here we study the problem of orienting a connected graph with cut edges in order to maximize the number of ordered vertex pairs  $(x,y)$  such that there is a directed path from  $x$  to  $y$ . After transforming this problem, we prove a key theorem about the transformed problem that allows us to obtain a quadratic algorithm for the original orientation problem. We also consider how to orient graphs to minimize the number of ordered vertex pairs joined by a directed path. After showing this problem is equivalent to the comparability graph completion problem, we show both problems are NP-hard, and even NP-hard to approximate to within a factor of  $1 + \epsilon$ , for some  $\epsilon > 0$ .

## 1 Introduction

Our terminology and notation is standard except as indicated. We mention only that if  $X$  is a subset of the vertices in a graph, we use  $\langle X \rangle$  to denote the subgraph induced by  $X$ . Good references for any other undefined terms are [1,2].

Let  $G$  be any connected graph. Given an orientation  $\vec{G}$  of  $G$ , we will use  $R(\vec{G})$  to denote the number of ordered vertex pairs  $(x, y)$  such that there is a directed path from  $x$  to  $y$  in  $\vec{G}$ . We call  $R(\vec{G})$  the *reachability* of  $\vec{G}$ .

Robbins [12] showed that  $G$  can be oriented so that  $\vec{G}$  is strongly connected (i.e.,  $R(\vec{G}) = |V|(|V| - 1)$ ) if and only if  $G$  is 2-edge-connected. In the following section (Section 2), we study the problem of how to orient an arbitrary graph  $G$  to obtain a digraph  $\vec{G}$  with  $R(\vec{G})$  as large as possible. We first transform the problem into an equivalent orientation problem on vertex-weighted trees. Although this transformed version is NP-hard, we prove a key theorem (Theorem 2.2) which allows us to get a quadratic algorithm for the original problem, as well as a fully polynomial approximation scheme for the transformed problem.

In Section 3, we consider the analogous problem of how to orient  $G$  so as to minimize  $R(\vec{G})$  for the resulting digraph  $\vec{G}$ . We show this problem is equivalent to comparability graph completion (adding the fewest edges so the resulting graph can be transitively oriented), and then show that both problems are NP-hard, and even NP-hard to approximate within a factor of  $1 + \epsilon$ , for some  $\epsilon > 0$ . Related hardness results appear in [6,10].

---

\*This work was supported by the National Science Foundation under Grants NCR-95-05551 and DMS-92-06991

<sup>†</sup>Department of Electrical and Computer Engineering, University of California, Davis, CA 95616

<sup>‡</sup>Department of Mathematics and Computer Science, San Jose State University, San Jose, CA 95192

<sup>§</sup>Department of Computer Science, Dartmouth College, Hanover, NH 03755

## 2 Orienting Graphs to Maximize Reachability

Suppose we are given a connected graph  $G$  on  $n$  vertices with cut edges. Our goal is to orient  $G$  to obtain a digraph  $\vec{G}$  with  $R(\vec{G})$  as large as possible.

It will be useful to first transform this basic problem. Let  $C_1, C_2, \dots, C_b$  denote the components left when the cut edges of  $G$  are removed from  $G$ . By Robbins Theorem [12] each  $C_i$  can be oriented into a strongly connected digraph. Let us contract each non-trivial  $C_i$  into a single vertex  $x_i$ , giving  $x_i$  weight  $wt(x_i) = |V(C_i)|$ . The resulting contracted graph is, of course, a  $b$ -vertex tree  $T = T(G)$  with integer weights on the vertices. Our original problem is now easily seen to be equivalent to the following problem: How should we orient  $T$  to maximize  $\sum wt(x_i) \cdot wt(x_j)$ , the sum being taken over all vertex pairs  $(x_i, x_j)$  in  $T$  such that there is a directed path from  $x_i$  to  $x_j$  in  $\vec{T}$ ? Note that the input size of the transformed problem may be exponentially smaller than the input size of the original problem (roughly,  $b(1 + \log n)$  versus  $n$ ).

We begin by showing that this transformed problem is NP-hard. In particular, consider the following decision problem:

### WEIGHTED TREE ORIENTATION (WTO)

**Instance:** Tree  $T$ , weight function  $wt: V(T) \rightarrow \mathbb{Z}^+$ , integer  $B > 0$ .

**Question:** Is there an orientation  $\vec{T}$  such that  $\mu(\vec{T}) \doteq \sum wt(v) \cdot wt(w) \geq B$ , the sum taken over all pairs  $(v, w)$  with a directed path from  $v$  to  $w$  in  $\vec{T}$ ?

**Theorem 2.1.** *WTO is NP-complete.*

**Proof.** WTO is obviously in NP, and so it suffices to show it is NP-hard. For this we reduce PARTITION [4,p.233] to WTO. Given positive integers  $a_1, a_2, \dots, a_m$  with even sum  $S$ , consider the weighted tree  $T$  in Fig. 1 and set  $B = 5(S/2)^2$ . Given any  $I \subseteq \{1, 2, \dots, m\}$ , consider the orientation  $\vec{T}$  obtained by orienting toward (resp, away from) the vertex with weight  $S$  each edge whose other end vertex has weight  $a_i$  for  $i \in I$  (resp,  $i \in \{1, 2, \dots, m\} - I = \bar{I}$ ). We then find


$$\mu(\vec{T}) = S \left( \sum_{i \in I} a_i + \sum_{i \in \bar{I}} a_i \right) + \sum_{i \in I} a_i \cdot \sum_{i \in \bar{I}} a_i \leq S^2 + \left( \frac{S}{2} \right)^2 = 5 \left( \frac{S}{2} \right)^2$$

with equality precisely if  $\sum_{i \in I} a_i = \sum_{i \in \bar{I}} a_i = \frac{S}{2}$ . ■

We now wish to develop a key result which will allow us to obtain a quadratic algorithm for the original orientation problem, as well as a fully polynomial approximation scheme for WTO. First, however, we need some terminology and notation. Let  $T$  be any tree with positive integer vertex weights. An *optimal orientation*  $\vec{T}$  is one which maximizes  $\mu(\vec{T})$ . Given an orientation  $\vec{T}$  and a vertex  $w \in V(T)$ , define

$$\text{In}_{\vec{T}}(w) \doteq \{x \in V(T) \mid \text{there exists a directed path from } x \text{ to } w \text{ in } \vec{T}\}$$

$$\text{Out}_{\vec{T}}(w) \doteq \{x \in V(T) \mid \text{there exists a directed path from } w \text{ to } x \text{ in } \vec{T}\}$$



**Figure 1.** Tree  $T(a_1, \dots, a_m)$

In particular,  $w \in \text{In}_{\vec{T}}(w) \cap \text{Out}_{\vec{T}}(w)$ . Define

$$\text{In}'_{\vec{T}}(w) \doteq \text{In}_{\vec{T}}(w) - \{w\} \text{ and } \text{Out}'_{\vec{T}}(w) \doteq \text{Out}_{\vec{T}}(w) - \{w\}.$$

We will usually drop the subscript  $\vec{T}$  if it is clear from the context. Finally, given any subgraph with vertex set  $X \subseteq V(T)$ , let  $\|X\| \doteq \sum_{x \in X} wt(x)$  be called the *total weight* of that subgraph. A *centroid* in a vertex weighted tree is any vertex  $c$  whose removal minimizes the maximum total weight of any component in  $T - c$ .

Our goal now is to prove the following.

**Theorem 2.2.** *Let  $c$  be a centroid of  $T$ . In every optimal orientation  $\vec{T}$ , we have  $\text{In}_{\vec{T}}(c) \cup \text{Out}_{\vec{T}}(c) = V(T)$ .*

Before proving this, we require the following result.


**Lemma.** *Let  $\vec{T}$  be any optimal orientation of  $T$ , and let  $P$  be any undirected path in  $T$ . Then there are no two vertices  $x, y$  on  $P$  such that the edges of  $P$  incident to  $x$  (resp,  $y$ ) are both directed toward  $x$  (resp, away from  $y$ ) in  $\vec{T}$ .*

Briefly, there are no "double reversals" on any path in  $\vec{T}$  (see Fig. 2).

**Proof of the Lemma.** Otherwise, let  $x, y$  be a closest such pair on  $P$ , so the edges of  $P$  between  $x$  and  $y$  form a directed path from  $y$  to  $x$ . Let  $e_x$  denote the edge at  $x$  on the path from  $y$  to  $x$ . Let  $X = \{w \in V(T) \mid \text{there is an undirected path in } T \text{ between } w \text{ and } x \text{ which does not contain } e_x\}$  and let  $X' = X \cap \text{In}(x)$ .

Let  $z$  denote the neighbor of  $y$  on  $P$  which does not occur on the directed path from  $y$  to  $x$  (see Fig. 2). Thinking of  $T$  as rooted at  $z$ , consider the subtree  $T_y$  rooted at  $y$ . Note that  $\|\text{In}'(y)\| \geq \|\text{Out}'(y) \cap T_y\|$  (else we could reverse the orientations of all edges in  $T_y$  to obtain an orientation better than  $\vec{T}$ ). Reverse all edges in  $\langle X' \rangle$ , and consider the gain and loss in  $\mu(\vec{T})$  in doing so, where the loss in  $\mu(\vec{T})$  is the sum of the terms in  $\mu(\vec{T})$  which no longer exist under the new orientation, and the gain in  $\mu(\vec{T})$  is defined analogously. Since  $\text{Out}'(x) \subset \text{Out}'(y) \cap T_y$ , we find

$$\begin{aligned} \text{Loss in } \mu(\vec{T}) &\leq \|X'\| \cdot \|\text{Out}'(x)\| < \|X'\| \cdot \|\text{Out}'(y) \cap T_y\| \\ &\leq \|X'\| \cdot \|\text{In}'(y)\| \leq \text{Gain in } \mu(\vec{T}) \end{aligned}$$



**Figure 2. Double Reversal on P**

and thus we would have a better than optimal orientation. This proves the Lemma.

**Proof of Theorem 2.2.** Throughout, think of  $T$  as rooted at centroid  $c$ . If the theorem fails for some optimal orientation  $\vec{T}$ , there must be a vertex  $x \notin \text{In}(c) \cup \text{Out}(c)$  adjacent to a vertex  $v \in \text{In}'(c) \cup \text{Out}'(c)$ . (We will call  $(v, x)$  a *dangling edge* at  $v$ .) We now consider two cases, assuming  $|\text{Out}(c)| \geq |\text{In}(c)|$  (else reverse the orientation of all edges of  $\vec{T}$ ).

**Case 1.** *There is a dangling edge  $(v, x)$  at  $v \in \text{In}'(c)$ .*

Note that since  $c \in \text{In}(c) - (T_v \cap \text{In}(c))$ , we have

$$|\text{Out}(c)| \geq |\text{In}(c)| > |T_v \cap \text{In}(c)|.$$

Reverse  $(v, x)$  and the edges in  $T_x$ . We find

$$\begin{aligned} \text{Gain in } \mu(\vec{T}) &\geq |\text{Out}(c)| \cdot |\text{Out}(x)| \\ &> |T_v \cap \text{In}(c)| \cdot |\text{Out}(x)| \geq \text{Loss in } \mu(\vec{T}) \end{aligned}$$

contradicting the optimality of  $\vec{T}$ .

**Case 2.** *There are no dangling edges at any vertex of  $\text{In}'(c)$ .*

Let  $(x, v)$  be a dangling edge at  $v \in \text{Out}'(c)$ . If  $\text{outdeg}(c) \geq 2$ , then  $\vec{T}$  contains a path which violates the Lemma (with  $v, c$  playing the roles of  $x, y$  in the Lemma). Hence we may assume  $\text{outdeg}(c) = 1$ . But then, since  $c$  is a centroid and there are no dangling edges off vertices in  $\text{In}'(c)$ , and since  $\{x\} \cup (T_v \cap \text{Out}(c)) \subset V(T) - \text{In}(c)$ , we find

$$|\text{In}(c)| \geq |V(T) - \text{In}(c)| > |T_v \cap \text{Out}(c)|.$$

Reverse  $(x, v)$  and the edges in  $T_x$ . We find

$$\text{Gain in } \mu(\vec{T}) \geq |\text{In}(c)| \cdot |\text{In}(x)| > |T_v \cap \text{Out}(c)| \cdot |\text{In}(x)| \geq \text{Loss in } \mu(\vec{T})$$

contradicting the optimality of  $\vec{T}$ . ■

It is well known that the centroid of a vertex-weighted tree can be found in linear time, and that the centroid consists of either a single vertex or two adjacent vertices [9]. In the latter case, Theorem 2.2 implies that the optimal orientation may be represented schematically as in Figure 3, where  $c_1$  and  $c_2$  denote the adjacent centroids.

In the former case, we have essentially an instance of PARTITION. Let  $T$  be rooted at  $c$ , and let  $v_1, \dots, v_k$  denote the neighbors of  $c$ . Consider an optimal orientation of  $\vec{T}$  of  $T$ . By Theorem 2.2, the subtree  $T_{v_i}$  must be oriented entirely toward (respectively, away from)  $c$  if and only if the edge between  $v_i$  and  $c$  is oriented toward (respectively, away from)  $c$ . Thus,




Figure 3.

$$\mu(\vec{T}) = \|c\| \cdot \|T - c\| + \left( \sum_{i \in I} \|T_{v_i}\| \right) \left( \sum_{i \in \bar{I}} \|T_{v_i}\| \right) + \sum_{i=1}^k \mu(T_{v_i}^*),$$

where  $I = \{i \mid T_{v_i} \text{ is oriented toward } c \text{ in } \vec{T}\}$  and  $T_{v_i}^*$  denotes  $T_{v_i}$  oriented entirely towards (respectively, away from)  $c$  if  $i \in I$  (respectively,  $i \in \bar{I}$ ). To maximize  $\mu(\vec{T})$ , we need to find a partition  $I \cup \bar{I}$  of  $\{1, 2, \dots, k\}$  such that  $\sum_{i \in I} \|T_{v_i}\|$  and  $\sum_{i \in \bar{I}} \|T_{v_i}\|$  are as equal as possible.

There is, of course, a well-known dynamic programming algorithm to find such a partition [4, Section 4.2]. Since  $\sum_{i=1}^k \|T_{v_i}\| \leq n$ , the running time of this dynamic programming algorithm is  $O(kn) = O(n^2)$ . Since all other tasks in the original orientation algorithm (e.g., finding the 2-edge-connected components in  $G$ , giving these components strongly-connected orientations, etc.) can easily be done in  $O(|E|)$  time using standard depth-first search techniques, we see that the original orientation problem can be solved in  $O(n^2)$  time; i.e., quadratic in the original input size  $n$ . On the other hand, there is also a well-known fully polynomial approximation scheme for the above partition problem [7], which in turn provides a fully polynomial approximation scheme for the NP-complete problem WTO.

### 3 Orienting Graphs to Minimize Reachability

Let  $G$  be a connected graph, and let  $r(G) = \min R(\vec{G})$ , the minimum being taken over all orientations of  $G$ . We call an orientation  $\vec{G}$  a *minimal orientation* if  $R(\vec{G}) = r(G)$ . We begin with the following result.

**Theorem 3.1.** *Every minimal orientation of  $G$  is acyclic.*

**Proof.** Suppose to the contrary there exists a minimal orientation  $\vec{G}$  which is not acyclic. Since  $\vec{G}$  is not acyclic, at least one of the strongly-connected components of  $\vec{G}$ , say  $C$ , is not a single vertex. Let  $E'$  denote a set of edges in  $C$  whose reversal renders  $\langle V(C) \rangle$  not strongly connected, and let  $\vec{G}'$  denote  $\vec{G}$  with the edges in  $E'$  reversed. It is easy to see every ordered pair of vertices joined by a directed path in  $\vec{G}'$  is joined by a directed path in  $\vec{G}$ , while clearly there is a pair of vertices in  $V(C)$  which are joined by a directed path in  $\vec{G}$  but not in  $\vec{G}'$ . Thus,  $R(\vec{G}') < R(\vec{G})$ , which violates the minimality of  $\vec{G}$ . ■

Obviously  $r(G) \geq |E(G)|$ , and it is known that  $r(G) = |E(G)|$  if and only if  $G$  is a comparability graph [1,5], or, equivalently, if  $G$  is transitively orientable [3]. It is known that comparability graphs can be recognized in polynomial time [3].

Given a connected graph  $G$ , let  $\underline{c}(G)$  (respectively,  $\bar{c}(G)$ ) denote the minimum number of edges which must be deleted from (respectively, added to)  $G$  to obtain a comparability graph. We now apply Theorem 3.1 to establish a connection between  $r(G)$  and  $\bar{c}(G)$ .

**Theorem 3.2.**  $r(G) = |E(G)| + \bar{c}(G)$ .

**Proof.** Suppose we add  $\bar{c}(G)$  edges to  $G$  to obtain a comparability graph  $G'$ , and then orient  $G'$  so that  $R(\vec{G}') = |E(G')| = |E(G)| + \bar{c}(G)$ . But, of course  $r(G) \leq R(\vec{G}')$ , and so  $r(G) \leq |E(G)| + \bar{c}(G)$ .

On the other hand, consider a minimal orientation  $\vec{G}$  of  $G$ , so that  $r(G) = R(\vec{G})$ . By Theorem 3.1,  $\vec{G}$  is acyclic. Consider the transitive closure  $cl(\vec{G})$  of  $\vec{G}$ . We then find

$$\begin{aligned} r(G) &= R(\vec{G}) = R(cl(\vec{G})) \\ &= |E(G)| + (\text{No. of Edges Added to } \vec{G} \text{ to obtain } cl(\vec{G})) \\ &\geq |E(G)| + (\text{Min. No. of Edges which need to be} \\ &\quad \text{added to } G \text{ to obtain a transitively orientable graph}) \\ &= |E(G)| + \bar{c}(G). \end{aligned}$$

Thus,  $r(G) = |E(G)| + \bar{c}(G)$ , as asserted. ■

We now turn to the complexity of computing  $r(G)$ . Consider the following decision problem.

### MINIMUM REACHABILITY ORIENTATION (MRO)

**Instance:** Graph  $G$ , integer  $k \geq |E(G)|$

**Question:** Is  $r(G) \leq k$ ?

In a moment we will prove

**Theorem 3.3.** *MRO is NP-complete.*

It follows immediately from Theorems 3.2 and 3.3 that the following problem is also NP-complete.

### COMPARABILITY GRAPH COMPLETION (CGC)

**Instance:** Graph  $G$ , integer  $k \geq 0$

**Question:** Is there a superset  $E'$  of  $E$  such that  $|E' - E| \leq k$  and  $G = (V, E')$  is a comparability graph (i.e., is  $\bar{c}(G) \leq k$ )?

(Previously, it was known only that COMPARABILITY SUBGRAPH (i.e., deciding if  $\underline{c}(G) \leq k$ ) is NP-complete [4,p.197]. For the optimization versions of MRO and CGC, the goal is to compute (or approximate) the minimum  $k$  such that  $\langle G, k \rangle$  is a positive instance of the decision problem.)

**Proof of Theorem 3.3.** Clearly  $\text{MRO} \in \text{NP}$ . To show MRO is NP-hard, we will reduce NOT-ALL-EQUAL 3SAT [4,p.259].

Let  $I$  be an instance of NAE3SAT with  $m$  clauses. Construct a graph  $G_I$  as follows. Each variable  $x$  will be represented by an edge  $(xT, xF)$  in  $G_I$ , and orienting this edge towards  $xT$  (respectively,  $xF$ ) will correspond to setting  $x$  to  $T$  (respectively,  $F$ ). Each clause  $C$  will be represented by a 9-cycle in  $G_I$ . The three literals in  $C$  will be assigned to three equally spaced edges  $(1T, 1F), (2T, 2F), (3T, 3F)$  on  $C$ 's 9-cycle, as shown in Fig. 3. If the literal assigned to the edge  $(1T, 1F)$  is  $X$  (respectively,  $\bar{X}$ ), add two 2-paths joining  $1T$  to  $xF$  and  $1F$  to  $xT$  (respectively, joining  $1T$  to  $xT$  and  $1F$  to  $xF$ ) to  $G_I$ . Make analogous connections for the other two literals of  $C$ , as well as for the remaining clauses of  $I$ , to complete  $G_I$ .




Figure 4. The Graph  $G_I$

Note that however we orient  $G_I$ , each of the  $m$  9-cycles will contain a directed 2-path, and thus  $r(G_I) \geq |E(G_I)| + m$ . In fact,  $r(G_I) = |E(G_I)| + m$  if and only if  $I$  has a satisfying truth assignment in the not-all-equal sense. Indeed, given a satisfying truth assignment for  $I$ , we can obtain such an orientation for  $G_I$  as follows: Orient the edges  $(xT, xF)$  to correspond to the truth assignment, and orient all the 6-cycles in  $G_I$  containing these edges so that none contains a directed 2-path. Note that the edges in each 9-cycle to which a literal was assigned are now oriented to reflect the truth value of that literal under the truth assignment. Since each clause contains both a true and a false literal, it is trivial to complete the orientation of the 9-cycles so each contains exactly one directed 2-path passing through a darkened vertex on the 9-cycle. Conversely, any orientation of  $G_I$  with only  $|E(G_I)| + m$  reachable pairs must have exactly one directed 2-path per 9-cycle, and so corresponds to a satisfying truth assignment. ■

We also observe that it remains NP-hard even to approximate  $r(G)$  to within a factor of  $1 + \epsilon$ , for some  $\epsilon > 0$ . Consider the following optimization problem.

### MAX NOT-ALL-EQUAL 3SAT

**Instance:** Boolean formula in 3CNF

**Question:** What is the maximum number of clauses that can be satisfied (in a not-all-equal sense) by a truth assignment?

It was established recently [8,11] that it is NP-hard to approximate MAX NAE3SAT to within a factor of 1.013 (that is, finding an assignment that satisfies, in the not-all-equal sense,  $1/1.013$  of the maximum possible number of clauses is NP-hard). Using this, we can strengthen Theorem 3.3 as follows:

**Theorem 3.4.** *There exists a constant  $\epsilon > 0$  such that approximating the optimization versions of MRO or CGC to within a factor of  $1 + \epsilon$  is NP-hard.*

**Proof Sketch.** Let  $I$  be an instance of NAE3SAT with  $m$  clauses. In the following, “satisfying” a clause of  $I$  refers to making at least one of its literals true and at least one false. Recall that  $R(\vec{G}_I)$  denotes the number of ordered pairs of vertices  $(x, y)$  with a directed path from  $x$  to  $y$  in an orientation  $\vec{G}_I$  of  $G_I$ .

**Claim 1:** *Given any assignment satisfying  $x$  of the clauses of  $I$ , there is an orientation  $\vec{G}_I$  of  $G_I$  such that  $R(\vec{G}_I) = |E(G_I)| + 3m - 2x$ .*

Claim 1 follows from the proof of Theorem 3.3.

**Claim 2:** *Given any orientation  $\vec{G}_I$ , there is an assignment satisfying at least  $\frac{1}{2}(|E(G_I)| + 3m - R(\vec{G}_I))$  clauses of  $I$ .* Claim 2 holds because if any clause-subgraph (see Fig. 4) in  $G_I$  has more than one directed 2-path, then it has at least three. Thus, any orientation can be converted into an equally good orientation corresponding to a truth assignment (where the only directed 2-paths are 2-paths through the darkened vertices on the 9-cycles). By simple algebra, the claimed bound holds for this assignment.

Suppose one could find an orientation  $\vec{G}_I$  of  $G_I$  with  $R(\vec{G}_I) \leq (1 + \epsilon)r(G_I)$ . By Claim 2, this would yield an assignment satisfying at least

$$(1) \quad \frac{1}{2}[|E(G_I)| + 3m - (1 + \epsilon)r(G_I)]$$

clauses of  $I$ . Let  $\max(I)$  denote the maximum number of clauses in  $I$  which can be simultaneously satisfied in a not-all-equal sense. By Claim 1,  $r(G_I) \leq |E(G_I)| + 3m - 2\max(I)$ . Thus expression (1) is at least  $\frac{1}{2}[|E(G_I)| + 3m - (1 + \epsilon)(|E(G_I)| + 3m - 2\max(I))]$ , which equals

$$(2) \quad \max(I) - \frac{\epsilon}{2}[|E(G_I)| + 3m - 2\max(I)].$$

Using  $|E(G_I)| \leq 24m$  and  $m \leq 2\max(I)$  (assuming without loss of generality that every clause has at least two literals, a random assignment satisfies at least half the clauses on average), expression (2) is at least  $\max(I) - \frac{\epsilon}{2}[48\max(I) + 6\max(I) - 2\max(I)] = (1 - 26\epsilon)\max(I)$ . From this the claimed hardness of approximating MRO follows, with  $\epsilon > 0.00049$ .

A similar argument, using Theorem 3.2 and omitting the  $|E(G_I)|$  term in expression (1) (and subsequent expressions), establishes the claimed hardness of approximating CGC, with  $\epsilon > 0.0064$ . ■



## 4 Concluding Remarks

It would be interesting to determine the algorithmic complexity of completing the orientation of a partially oriented graph to maximize reachability. Thus far we have made little progress on this problem. The analogous completion problem to minimize reachability is, of course, NP-hard.

On the other hand, it is very easy to characterize partial orientations which can be completed into strong orientations. An edge-cut  $(X, \overline{X})$  in a partially oriented graph is called *one-way* if all the edges in the cut are already oriented, and all are oriented from  $X$  to  $\overline{X}$  or all from  $\overline{X}$  to  $X$ . We have

**Theorem 4.1.** *A partial orientation of a graph  $G$  can be completed into a strong orientation if and only if  $G$  is 2-edge-connected and there are no one-way edge cuts in the partial orientation.*

## 5 References

- [1] C. Berge, *Graphs and Hypergraphs*, North-Holland, Amsterdam, 1973.
- [2] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications*, North-Holland, Amsterdam, 1976.
- [3] S. Even, A. Pnueli, and A. Lempel, "Permutation graphs and transitive graphs," *J. Assoc. Comp. Mach.* **19** (1972), 400-410.
- [4] M.R. Garey and D.S. Johnson, *Computers and Intractability: A Guide to NP-Completeness*, W.H. Freeman, San Francisco, 1979.
- [5] P.C. Gilmore and A.J. Hoffman, "A characterization of comparability graphs and of interval graphs," *Can. J. Math.* **16** (1964), 539-548.
- [6] M.C. Golumbic, H. Kaplan and R. Shamir, "Graph sandwich problems," *J. Algorithms* **19** (1995), 449-473.
- [7] O.H. Ibarra and C.E. Kim, "Fast approximation algorithms for the knapsack and sum of subset problems," *J. Assoc. Comp. Mach.* **22** (1975), 463-468.
- [8] V. Kann, J. Lagergren, and A. Panconesi, "Approximability of maximum splitting of  $k$ -sets and some other APX-complete problems," *Tech. Report TRITA-NA-P9509*, Dept. of Numer. Analysis and Comp. Sci., Royal Inst. of Technology, Stockholm.
- [9] O. Kariv and S.L. Hakimi, "An algorithmic approach to network location problems, II: the  $p$ -medians," *SIAM J. Appl. Math.* **37**(1979), 539-560.
- [10] P. Dell'Olmo, M. Grazia Speranza and Z. Tuza, "Comparability graph augmentation for some multiprocessor scheduling problems," *Discrete Appl. Math.* **72** (1997), 71-84.
- [11] C.H. Papadimitriou and M. Yannakakis, "Optimization, approximation, and complexity classes," *J. Comp. Systems Sci.*, **43** (1991), 425-440.
- [12] H.E. Robbins, "A theorem on graphs with an application to a problem of traffic control," *Amer. Math. Monthly* **46** (1939), 281-283.